

## Urea as the Nitrogen Source in NFT Hydroponic System

AMINUDDIN H., KHALIP R.<sup>1</sup>, NORAYAH K. and ALIAS H.<sup>2</sup>

Department of Soil Science, Universiti Pertanian Malaysia

<sup>1</sup>Agronomy Department, Universiti Pertanian Malaysia

<sup>2</sup>Petroleum Research Institute, PETRONAS

**Keywords:** Hydroponics, urea, hydrolysis, urease, ammonium

### ABSTRAK

Unsur nitrogen dalam larutan nutrien hidroponik cara NFT (nutrient film technique) menjalani proses hidrolisis dengan pengeluaran ammonium. Hidrolisis urea berlaku dengan pesat dari hari ketujuh dan tamat pada hari kedua puluh. Dalam masa yang sama jumlah ammonium dalam larutan meningkat dan mencapai tahap maksima pada hari kedua puluh. Penitratan berlaku serentak dan kepekatan nitrat mencapai maksima pada hari kedua puluh juga. Berat kering tanaman pada peringkat matang tidak berbeza antara larutan rawatan urea dan rawatan nitrat. Kepekatan ammonium dalam daun dan akar rawatan urea adalah tinggi dari rawatan nitrat. Kepekatan nitrat dalam daun dan akar tidak berbeza diantara kedua dua rawatan tersebut. Kajian menunjukkan urea dapat menggantikan nitrat sebagai sumber unsur nitrogen untuk tanaman dalam sistem hidroponik cara NFT.

### ABSTRACT

Urea as the source of nitrogen in the nutrient solution in NFT (nutrient film technique) hydroponic system undergoes hydrolysis which results in the release of ammonium in solution. Urea hydrolysis was rapid from the 7th day onwards and ended by the 20th day. At the same time, ammonium concentration in solution increased and reached its maximum on the 20th day. Nitrification occurred simultaneously and peaked also on the 20th day. Plant dry matter weight at harvest was similar for both urea and nitrate treatments. Ammonium concentration in leaves and roots was higher in urea than in nitrate treatments. Nitrate concentration in leaves and roots was similar for both treatments. The study showed that urea can be substituted for nitrate as the nitrogen source in the NFT hydroponic system.

### INTRODUCTION

The source of nitrogen in the nutrient solution for the growing of crops under the hydroponic system has been in the forms of  $\text{Ca}(\text{NO}_3)_2$  and  $\text{KNO}_3$  (Cooper 1975). Even though urea is increasingly being used as the main source of nitrogen fertilizer for crops grown on soil (Gould *et al.* 1986), its use as the N source for crops grown under the hydroponic system has yet to be evaluated. One of the reasons which could hinder the use of urea in hydroponics is that urea has to undergo hydrolysis with the release of  $\text{NH}_4^+$  and subsequent formation of  $\text{NO}_3^-$  through nitrification, forms that are utilised by the plants. Hydrolysis of urea proceeds through enzymatic catalysis of urease (Vlek *et al.* 1980). In soils urease activity is located in the soil biomass and solution originating from decaying

organisms (Skujins 1976; Vlek *et al.* 1980). There is also evidence that soil urease activity may be derived from plants (Frakenberger and Tabatabai 1982). It has also been shown that floodwater of tropical lowland rice soils has measurable amounts of urease activity (Sahrawat 1980), which could possibly originate from the soil.

Since the presence of urease has always been associated with soil, its presence in the nutrient solution in the hydroponic system devoid of soil therefore is of interest. Since bacteria can proliferate in nutrient solution and urease originates from bacteria cells, the possible presence of urease in nutrient solution media and hydrolysis of urea in the solution therefore is great.

The objective of this study was to examine the possibility of using urea as the nitrogen source in the nutrient solution in the hydroponic system by following its transformation in the nutrient solution and the performance of *Brassica chinensis* grown in it.

### MATERIALS AND METHODS

The NFT (nutrient film technique) of soilless culture with plants grown in channels fed with recirculating nutrient solution was used (Cooper, 1975). The container which held the recirculating nutrient solution had a capacity of 50 l. The channels were 3.7 m long, 30 cm wide and 5 cm high. These channels were constructed from 1 cm thick plywood. The inner surface of the channels was covered with polythene sheets along which the solution flowed and bathed the plant roots. A submersible pump in the container delivered the solution through 2 cm (diameter) PVC pipes from the container to the top of the channel, discharging the solution through a 5 mm tube at the rate of 2 l/min. The channels were elevated at the top end (10 cm higher) so that the solution would flow in a thin film on the surface of the channel to the bottom end and into the container and then recirculated. Each replicate of a treatment consisted of the channel, a container and a pump. Three replicates were used for each treatment. The experiment was run in the greenhouse in a complete randomized design. Mini pak choy (*Brassica chinensis* var Ching Chiang -Taiwan) seeds were sown on rockwool blocks 6 cm<sup>2</sup> and 40 cm in length for each block; nine blocks were used for each channel. These blocks were placed at the centre of the channels. Three seeds were placed at each point, 8 cm apart. Each rockwool block had 5 plants. Thinning was done on the tenth day leaving one plant per point. The total number of plants per channel at the start of the experiment was 45. Since the concentration of urea in the solution had to be monitored continuously for the whole period of the plant growth, elemental nutrients were added to the solution for plant need, based on the electrical conductivity (E.C.) of the solution; the element content of the solution was analysed on the 20th day to ensure that all elements with the exception of N were adequate for plant need. Plants were observed for deficiency symptoms. Cooper and Charlesworth (1977) concluded that addition of nutrients under the NFT based on the

readings of the E.C. and the solution pH was adequate to produce equivalent tomato yield compared to nutrient addition based on nutrient analysis.

### Treatments

The standard nutrient formulation used was based on Cooper (1975) (Table 1). The other three treatments were 100% N-urea (No.2), mixture of 50%N-nitrate and 50%N-urea (No.3) and 75%N-urea and 25%N-nitrate (No.4). An accompanying treatment (No.5) similar to (No.2) but without plants grown was included. CaCl<sub>2</sub>, KOH and K<sub>2</sub>SO<sub>4</sub> were substituted for the Ca(NO<sub>3</sub>)<sub>2</sub> and KNO<sub>3</sub> for treatments containing urea, resulting in some treatments having higher sulphur and chloride contents which, however, did not exceed toxic levels. pH of the solution was maintained at between 5.5-6.5 by adding HCl or NaOH. Electrical conductivity (E.C.) of the solution was maintained at 2.4 dS m<sup>-1</sup> until the plants were harvested. Solution E.C. which fell below this value was replenished with the stock solution of the respective treatments. Amounts added to each treatment are given in Table 2. Water loss through evapotranspiration was replaced with tap water to the 50 l mark daily.

### Sampling

Solution samples were taken daily for the first seven days and thereafter on the 10th, 20th,

TABLE 1  
Chemical make-up of the various treatments

Source	Treatments (g/1000)				
	1	2	3	4	5
urea	-	480	240	360	480
KH <sub>2</sub> PO <sub>4</sub>	272	272	272	272	272
KNO <sub>3</sub>	808	-	808	404	-
Ca(NO <sub>3</sub> ) <sub>2</sub>	943	-	-	-	-
CaCl <sub>2</sub>	-	588	588	588	588
MgSO <sub>4</sub>	492	492	492	492	492
KOH	-	224	-	112	224
K <sub>2</sub> SO <sub>4</sub>	-	348	-	174	348
MnSO <sub>4</sub>	5.27	5.27	5.27	5.27	5.27
H <sub>3</sub> BO <sub>3</sub>	3.03	3.03	3.03	3.03	3.03
Na <sub>2</sub> MoO <sub>4</sub>	.027	.027	.027	.027	.027
ZnSO <sub>4</sub>	.22	.22	.22	.22	.22
CuSO <sub>4</sub>	.05	.05	.05	.05	.05
EDTA-Fe	19.48	19.48	19.48	19.48	19.48

TABLE 2  
Nitrogen supplement during the period of  
plant growth

Treatments	NO <sub>3</sub> -N (g)	Urea-N (g)
T1	14.79	-
T2	-	6
T3	19.78	8.7
T4	7.38	10.8
T5	-	-

30th and the 40th day. Plants were harvested on the 40th day. PMA (phenylmercuric acetate) was added to samples to retard hydrolysis of urea before analysis of the samples was done. Samples were analysed for NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> (Bremner 1965) and urea (Douglas and Bremner 1970).

Plant samples were taken on the 20th, 30th and the 40th day. For the first two samplings, 9 plants (one from each rockwool block taken at random) were harvested from each channel. The tops and roots were separated and weighed. The final harvest comprised 27 plants. They were oven dried, reweighed and ground. For analysis of NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> in the samples, 1 g sample (oven dry) was shaken with 50 ml distilled water for 1 hr and the extract filtered. 10 ml of the extract was determined for NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> (Woolley *et al.* 1960).

## RESULTS AND DISCUSSION

Results of the analysis of the nutrient solution on the 20th day (Table 3) indicated that the macronutrient element content was adequate (Cooper and Charlesworth 1977). The elements content in the leaves at harvest also indicated that their concentration in plants were adequate (Table 3). Visual observation of the plants indicated no apparent nutrient deficiency except that the 100% N-urea treatment had darker green leaves than those in the other treatments.

Urea concentration in solution of treatments containing urea dropped rapidly from the 7th day onwards which showed that it had been hydrolysed. By the 20th day all the urea was depleted from the nutrient solution (Fig. 1). In soils urea is found to be fully hydrolysed within 3 days after application mostly in the form of NH<sub>4</sub><sup>+</sup> and by the 14th day most are in the NO<sub>3</sub><sup>-</sup> form (Gasser, 1964; Bundy and Bremner 1974). Decrease in the urea concentration of

the solution in treatments containing urea was rapid from the 7th day onwards. Loss of urea in the 100% N- urea with plant treatment was greater compared to the same treatment without plants on the 10th day. It is possible that the urease activity was greater in solution where plants were present. Elliott (1986) found hydrolysis of urea in cropped media more rapid than in uncropped media. Between the 1st and 7th day the urea loss was about 5%. Between the 7th and 10th day, urea in solution of cropped treatments decreased by 30%. Most of the urea was hydrolysed between the 10th and 20th day. Ammonium in the solution was detected as early as the first day after the start of the experiment in all the treatments receiving urea, indicating that hydrolysis of urea occurred immediately.




Fig. 1: Urea concentration in solution of the various treatments

### Ammonium in Solution

Ammonium concentration in the nutrient solutions increased until the 20th day, thereafter its concentrations began to decline except in the treatment without plants (Fig. 2). Plant uptake and the transformation of NH<sub>4</sub><sup>+</sup> to NO<sub>3</sub><sup>-</sup> via nitrification are the possible reasons for the decline after the 20th day. Furthermore, its source, urea, had been depleted by that time. Ammonium concentration in solution depended on the amount of urea in the treatments. High urea levels resulted in high NH<sub>4</sub><sup>+</sup> concentrations. The highest NH<sub>4</sub><sup>+</sup> concentration was recorded in the 100% N-urea treatments. With plant growth the decrease was higher which could be due to plant uptake and increased nitrification. Decline in NH<sub>4</sub><sup>+</sup> concentration in the treatment without plants after the 20th day

was slight. 75% and 50% N- urea treatments attained maximum  $\text{NH}_4^+$  concentration on the 20th day and the  $\text{NH}_4^+$  concentrations were in proportion to the amount of urea added. It can therefore be said that rate of hydrolysis of urea was not affected by the amount of urea present within the amounts of urea used in the experiment. This showed that urease was a non-limiting factor. Several researchers have found that in soils the rate of urea hydrolysed by soil urease increased with increase in urea concentration until the amount of urea added is sufficient to saturate the enzyme with the substrate (Douglas and Bremner 1971; Tabatabai and Bremner 1972; Dalal 1975).




Fig. 2:  $\text{NH}_4^+$  concentration in solution of the various treatments

*Nitrate in Solution*

The extent of nitrification occurring in the solution containing urea is shown in Fig. 3. Similar to the  $\text{NH}_4^+$  buildup,  $\text{NO}_3^-$  increase in the nutrient solution was observed in the 100% N-urea treatments with and without plants; however the buildup started later (from the 7th day onwards). While treatments with 100% and 50% N- $\text{NO}_3^-$  source declined for the period of 7- 10th day, treatments with 100%N-urea had  $\text{NO}_3^-$  concentration increased during the same period.

Maximum concentration of  $\text{NO}_3^-$  in the nutrient solution in the 100% N-urea treatments was attained on the 20th day. Subsequent periods showed no increase in nitrate concentration as shown by the 100% N- urea treatment without plants. At the 20th day the concentration of  $\text{NH}_4^+$  was only one third of  $\text{NO}_3^-$ .

Ammonium buildup and nitrate formation in the treatments containing urea occurred simultaneously. While 100% N-urea treatment with

plants had a  $\text{NO}_3^-$  concentration decrease after the 20th day, treatment without plants maintained the  $\text{NO}_3^-$  concentration until the 40th day. Nitrate concentration in treatments which contained urea decreased more sharply until the 40th day compared to plants in the 100% N- $\text{NO}_3^-$  treatment.




Fig. 3:  $\text{NO}_3^-$  concentration in solution of the various treatments

*Total N in solution*

Summation of the various forms of N ( $\text{NO}_3^-$ ,  $\text{NH}_4^+$  and urea) in the solution of the treatments is presented in Fig. 4. Total N remained the same for the first 7 days. From the 7th to 10th day total N of treatments 100% and 50% N- $\text{NO}_3^-$  was lower compared to the others. From the 10th to the 20th day , 100% and 75 % N- urea treatments decreased in total N value. Further decline in total N occurred in all treatments except the one without plants from the 20th to the 30th day. At the 40th day, treatments which contained urea had lower values compared to 100% N-  $\text{NO}_3^-$  treatment. From the 20th day onwards the treatment without plants maintained its total N to the end.




Fig. 4: Total N in solution of the various treatments

TABLE 3

Macronutrient concentrations in the solution (ppm) on the 20th. day and in the leaves at harvest (%)

Elements	Treatments			
	1	2	3	4
K (solution-s) (leaves-l)	306	332	249	263
	11.5	8.5	8.2	8.4
P (s) (l)	29	52	32	39
	1.33	1.10	1.04	1.22
Ca (s) (l)	175	214	227	200
	3.70	3.08	3.09	3.25
Mg (s) (l)	68	42	43	42
	.63	.56	.43	.50
N (l)	4.97	5.51	5.18	4.4

#### Plant dry weight

Plant dry weight on the 20th day for 100% N-urea treatment was the lowest recorded and was significantly different ( $P < 0.05$ ) from those of 100% and 50% N-nitrate treatments (Table 4). This initial difference in the dry weight can be attributed to the time taken for urea to be hydrolysed before  $\text{NH}_4^+$  and the subsequent release of  $\text{NO}_3^-$  in the solution for plant use. On the 30th day, the plant dry matter weight of the 100% N-nitrate treatment was not significantly different from that of 100%N-urea treatment. However, the two treatments with mixtures of urea and nitrate registered the highest value. Studies by Cox and Reisenauer (1973) have shown that mixtures of  $\text{NH}_4^+$  and  $\text{NO}_3^-$  in solution give higher yield than those of  $\text{NO}_3^-$  or  $\text{NH}_4^+$  supplied singly. Moneerat *et al.* (1982) reported that maximum dry weight was achieved with a ratio of 60: 40 ( $\text{NH}_4^+ : \text{NO}_3^-$ ) in the solution. Dry matter weight on the 40th day showed no significant difference among the four treatments (Table 4).

#### Total N Uptake by Plants

Total N uptake values normally follow dry matter weight figures. Total N uptake values of *Brassica* at harvest (40 days) were not significantly different among treatments. The 50% N-urea and 50% N-nitrate mixture recorded the highest N uptake value of 3.16 g. Baker and Maynard (1972) reported that maximum N ab-

sorption by plants was achieved when  $\text{NH}_4^+$  and  $\text{NO}_3^-$  were combined in the nutrient solution. The amount of N taken up by the plants accounts for 80% of the N supplied in the solution.

#### Ammonium in Roots

The percent coefficient of variation (%C.V.) of the  $\text{NH}_4^+$  content in the plant roots on the 20th day growth stage was high due to the problem of separating the roots from the rockwool media. At this stage the roots were very fine and were firmly embedded in the media making the task of separation difficult. At the 30th and 40th day, the roots were thicker and bigger and the rockwool was softer making the task of separation easier. This was reflected in the decreasing trend in the C.V. value as the plants matured. No significant difference in ammonium concentration in the plant roots of the 20- and 30-day-old plants of the various treatments was observed (Table 4). Plants treated with 100% N-nitrate did contain  $\text{NH}_4^+$  in the roots. The reduction and assimilation of  $\text{NO}_3^-$  in the roots produce  $\text{NH}_4^+$  as an intermediary product (Pate 1973).  $\text{NH}_4^+$  concentration in the roots of the 30- and 40-day old plants treated with 100%N-urea was higher than those treated with 100%N-nitrate. Pill and Lambert (1977) noted that in general, plants supplied with  $\text{NH}_4^+$  have a higher concentration of free  $\text{NH}_4^+$  than plants grown at comparable levels of  $\text{NO}_3^-$ .  $\text{NH}_4^+$  accumulation in the roots of the 100% N-urea treatment could also be due to the slow conversion of  $\text{NH}_3$  into glutamine which is a function of the carbohydrate supply (Vickery *et al.* 1936).

#### Ammonium in Leaves

Ammonium concentration in the leaves of 20-day-old plants treated with 100% N-urea was higher than that in plants in the 100% N-nitrate treatments (Table 4). Pate and Wallace (1964) found that the bulk of the  $\text{NH}_4^+$  absorbed by roots is converted to amino acids by the metabolic system of the roots and only a small percentage of the  $\text{NH}_4^+$  is found in the xylem exudate presumably transported to the leaves for assimilation. It is possible that in leafy vegetables, the leaves play a major role in the assimilation of  $\text{NH}_4^+$  which thus results in concentration of  $\text{NH}_4^+$  in leaves being comparable to that in the roots. On the 30th day,  $\text{NH}_4^+$  concentration in the leaves in all the treatments was similar.

TABLE 4  
Dry matter yield, total N,  $\text{NH}_4^+$  and  $\text{NO}_3^-$  content of pak choy

Dry matter yield (g/9 plants)		C.V. (%)	Treatments			
			1	2	3	4
D	20	21.89	1.26 a	0.75 b	1.26 a	1.21 ab
a	30	30.70	16.5 ab	10.77 b	21.61 a	19.43 ab
y	40	19.95	135.18 a	147.69 a	184.91 a	187.38 a
Total N uptake (g/9 plants)						
Day	40	23.86	2.20 a	2.75 a	3.16 a	2.76 a
$\text{NH}_4$ (roots) ppm						
D	20	91.58	824.6 a	803.6 a	200.0 a	304.3 a
a	30	29.72	570.0 b	1247.8 a	1470.0 a	1320.0 a
y	40	14.92	2010.0 b	2910.0 a	2610.0 ab	2610.0 ab
$\text{NH}_4$ (leaves) ppm						
D	20	43.66	324.5 b	693.8 a	330.0 b	360.0 ab
a	30	28.13	1260.0 a	1620.0 a	1680.0 a	1350.0 a
y	40	23.79	1680.0 b	2790.0 a	2100.0 ab	2340.0 ab
$\text{NO}_3$ (roots) %						
D	20	54.60	2.13 a	1.10 ab	0.90 ab	0.73 b
a	30	47.99	0.30 a	0.40 a	0.36 a	0.20 a
y	40	46.41	0.10 a	0.09 a	0.12 a	0.10 a
$\text{NO}_3$ (leaves) %						
D	20	46.30	3.27 a	2.27 a	2.73 a	2.38 a
a	30	33.78	2.48 a	1.63 ab	1.55 ab	1.21 b
y	40	46.93	1.01 a	0.99 a	1.21 a	0.89 a

Means with the same letter in a row are not significantly different at 0.05 using DMRT.

This period coincides with the high concentration of  $\text{NH}_4^+$  in the nutrient media. As the concentration of  $\text{NH}_4^+$  in solution started to fall by the 40th day,  $\text{NH}_4^+$  concentration in leaves of 100%N-nitrate plants decreased. Since 100% N-nitrate treatment plants had less than 5%  $\text{NH}_4^+$  content in the nutrient solution, we can deduce that the  $\text{NH}_4^+$  present in the the leaves was from the reduction of  $\text{NO}_3^-$ . The  $\text{NH}_4^+$  concentration in the leaves increased as the plant matured.

#### Nitrate in Roots

Nitrate concentration in roots was in contrast with that of  $\text{NH}_4^+$ . Whilst the concentration of

$\text{NH}_4^+$  increased with plant growth,  $\text{NO}_3^-$  concentration decreased.  $\text{NO}_3^-$ :  $\text{NH}_4^+$  concentration ratio was >20 in the 20-day-old plants and was <1 in the 40-day-old plants. Increased  $\text{NO}_3^-$  assimilation to amino acids in plants as they mature could be the reason for the fall in  $\text{NO}_3^-$  concentration. No significant difference in % nitrate in roots of plants treated with 100% N-nitrate and 100% N-urea at the 20th, 30th and 40th day was observed.

#### Nitrate in Leaves

$\text{NO}_3^-$  concentration in the leaves of pak choy was higher than in the roots. Pate(1973) found that

the reduction of  $\text{NO}_3^-$  concentration in the leaves is relative to the roots and varies widely among plant species. It is possible that  $\text{NO}_3^-$  assimilation in the leaves of pak choy is more intense than in the roots. In maize plants, the roots were found to reduce about 1/3 of the nitrate and the percentage of  $\text{NO}_3^-$  reduction decreased as the plant aged (Raghuveer 1977). No significant difference in % nitrate in leaves between treatments was observed at the three plant growth stages (Table 4). The amount of  $\text{NO}_3^-$  in the leaves decreased as the plant growth increased. Percent  $\text{NO}_3^-$  was ten-fold higher than  $\text{NH}_4^+$  in the leaves, the ratio being greater when plants were young.

### CONCLUSION

Urea as the source of N used in nutrient solution in the NFT hydroponic system undergoes hydrolysis with the release of  $\text{NH}_4^+$  and the subsequent formation of  $\text{NO}_3^-$ . The urea was completely hydrolysed by the 20th day. There was no difference in either the plant dry weight in the 100% N-urea or 100% N-nitrate treatments at harvest.  $\text{NH}_4^+$  concentration in roots and leaves was higher in the 100% N-urea treatment than in the 100% N-nitrate treatment. No difference in  $\text{NO}_3^-$  concentration in roots and leaves of 100% N-nitrate and 100% N-urea treatments was detected. The study showed the possibility of using urea as the N source for the NFT hydroponic system in crop production.

### REFERENCES

- BAKER, A.V and D.N. MAYNARD. 1972. Cation and nitrate accumulation in pea and cucumber plants as influenced by N nutrition. *J. Amer. Soc. Hort. Sci.* **97**: 27-30.
- BREMNER, J.M. 1965. Inorganic Forms of Nitrogen. In C.A. Black (ed.). *Methods of Soil Analysis, Part 2*. 1179-1232. Madison, Wisconsin: Am. Soc. Agron.
- BUNDY, L.G. and J.M. BREMNER. 1974. Effects of nitrification inhibitors on transformation of urea N in soils. *Soil Biochem.* **6**: 369-376.
- COOPER, A.J. 1975. Crop production in recirculating nutrient solution. *Scientia Hort.* **3**: 251-258.
- COOPER, A.J. and R.R. CHARLESWORTH. 1977. Nutritional control of a nutrient-film tomato crop. *Scientia Hort.* **7**: 189-195.
- COX, W.J. and H.M. REISENAUER. 1973. Growth and ion uptake by wheat supplied nitrogen as nitrate, or ammonium, or both. *Plant Soil* **38**: 363-380.
- DALAL, R.C. 1975. Urease activity in some Trinidad soils. *Soil Biol. Biochem.* **7**: 5-8.
- DOUGLAS, C.A. and J.M. BREMNER. 1970. Extraction and colorimetric determination of urea. *Soil Sci. Soc. Am. Proc.* **34**: 859-862.
- DOUGLAS, C.A. and J.M. BREMNER. 1971. A rapid method of evaluating different compounds or inhibitors of urease activity in soils. *Soil Biol. Biochem.* **3**: 309-315.
- ELLIOTT, G.C. 1986. Urea hydrolysis in potting media. *J. Am. Soc. Hort. Sci.* **6**: 862-866.
- FRAKENBERGER, W. T. and M.A. TABATABAI. 1982. Amidase and urease activities in plants. *Plant and Soil* **64**: 153-166.
- GASSER, J.K.R. 1964. Urea as a fertilizer. *Soils Fert.* **27**: 175-180.
- GOULD, W.D., C. HAGEDORN and R.G.L. MCCREADY. 1986. Urea transformations and fertilizer efficiency in soil. *Adv. in Agron.* **40**: 209.
- MONERAT, P.H., P.D. CASTALLANE, J.L.C. ZAMBON, J.G. PANDUA and J.J.V. MULLER. 1982. Effect of ammonium to nitrate ratio on initial development and mineral composition of tomato, *Lycopersicon esculentum*. *Proceedings of Amer. Soc. Hort. Sci.* **25**: 451-455.
- PATE, J.S. 1973. Uptake, assimilation and transport of nitrogen compounds by plants. *Soil Biol. Biochem.* **5**: 109-119.
- PATE, J.S. and W. WALLACE. 1964. Movement of assimilated nitrogen from the root systems of the field pea (*Pisum arvense* L.) *Ann. Bot.* **28**: 83-99.
- PILL, W.G. and V.N. LAMBERT. 1977. Effects of  $\text{NH}_4$  and  $\text{NO}_3$  nutrition with and without pH adjustments on tomato growth, ion compositions and water relations. *J. Am. Soc. Hort. Sci.* **102**: 78-81.
- RAGHUVEER, P. 1977. Characteristics of nitrate uptake from nutrient solution and root nitrate reductase activity among corn genotype. Ph.D Dissertation. Univ. of Illinois, Urbana-Champaign.

- SAHRAWAT, K.L. 1980. Urease activity in tropical rice soils and floodwater. *Soil Biol. Biochem.* **12**: 195-196.
- SKUJINS, J.J. 1976. Extracellular enzymes in soils. *CRC Crit. Rev. Microbiol.* **4**: 383-421.
- TABATABAI, M.A and J.M.BREMNER. 1972. Assay of urease activity in soils. *Soil Biol. Biochem.* **4**: 479-487.
- VICKERY, H.B., G.W. PUCHEZ and H.E. CLARK. 1936. Glutamine metabolism of the beet. *Plant Physiol.* **11**: 413-420.
- VLEK, P.L.G., J.M. STUMPE and B.H. BYRNES. 1980. Urease activity and inhibition in flooded soil systems. *Fert. Res.* **1**: 191-202.
- WOOLLEY, J.T., G.P. HICKS and R.H. HAGEMAN. 1960. Rapid determination of nitrate and nitrite in plant material. *J. Agric. and Food Chem.* **8**: 481-482.

(Received 27 August 1991)